



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

governor may also be cited. The whirligig may be considered a practical reversion of the principle exemplified in this problem.

AVERAGE AND PROBABILITY.

NOTE ON THE POND PROBLEM.

It was our purpose to publish a second solution of problem 90, Average and Probability, by a different method, but so far we have been unable to obtain a positive result. We have gone over our investigations a number of times, and have had several of our contributors go over the calculations, and so far we have been unable to find our error. Since we obtain a negative result, there is certainly something wrong with the work, for the problem admits of a definite solution when once the law of distribution has been decided upon. We hope to find time to go over our work again, and should we find our error, we will publish our solution as the method may be of interest to many of our readers.

This problem was originally proposed by Artemas Martin, Ph. D., and published as problem 300, Vol. I., No. 6, page 195, in the *Mathematical Visitor*, edited and published by himself. We infer that this is the source from which Dr. Byerly took it for his problem 21, *Integral Calculus*, second edition, page 211. While no specific reference is made as to the source from which it was taken, yet in his introductory paragraph on the subject of Mean Value and Probability, Dr. Byerly makes mention of the *Mathematical Visitor*.

In a letter to us from Dr. Martin, he says that Professor Seitz sent him a solution, giving the same answer as was obtained in the two solutions published in the December number of Vol. VII. ED. F.

106. Proposed by L. C. WALKER, A. M., Petaluma High School, Petaluma, Cal.

Required the average distance between two points in a hemisphere.

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Let O be the center of the sphere, P any point in the radius AO or BO . Take the left hemisphere when $OP < \frac{1}{2}r$, the right hemisphere when $OP > \frac{1}{2}r$. Let $OP = x$, PC , PG , PD , PE , $PF = y$.

If the first point is anywhere on the hemisphere with radius x , the second point must be on the surface of the zone of a sphere radius y . The surface of the hemisphere radius x is $2\pi x^2$ when $x < \frac{1}{2}r$. If $y < x$ the second point lies on the surface of a sphere radius y . The area of this surface $= 4\pi y^2$. If $y > x$ and $< (r-x)$, the second point is on the area $2\pi PR \cdot OR =$

